

Principal Component Analysis

Applied Multivariate Statistics – Spring 2012



Overview


- Intuition
- Four definitions
- Practical examples
- Mathematical example
- Case study

PCA: Goals

- Goal 1: Dimension reduction to a few dimensions
(use first few PC's)
- Goal 2: Find one-dimensional index that separates objects best
(use first PC)

PCA: Intuition


- Find low-dimensional projection with largest spread



PCA: Intuition




PCA: Intuition




PCA: Intuition

	X_1	X_2
Std. Basis	0.3	0.5
PC Basis	0.7	0.1
After Dim. Reduction	0.7	-




PCA: Intuition in 1d



Taken from “The Elements of Stat. Learning”, T. Hastie et.al.

PCA: Intuition in 2d



Taken from “The Elements of Stat. Learning”, T. Hastie et.al.


PCA: Four equivalent definitions

- Always center data first ! **Good for intuition**
- Orthogonal directions with largest variance
- Linear subspace (straight line, plane, etc.) with minimal squared residuals
- Using Spectraldecompsition (=Eigendecomposition)
- Using Singular Value Decomposition (SVD)

Good for computing


PCA (Version 1): Orthogonal directions

- PC 1 is direction of largest variance
- PC 2 is
 - perpendicular to PC 1
 - again largest variance
- PC 3 is
 - perpendicular to PC 1, PC 2
 - again largest variance
- etc.



PCA (Version 2): Best linear subspace

- PC 1: Straight line with smallest orthogonal distance to all points
- PC 1 & PC 2: Plane with with smallest orthogonal distance to all points
- etc.



PCA (Version 3): Eigendecomposition

- **Spectral Decomposition Theorem:**

Every symmetric, positive semidefinite Matrix R can be rewritten as

$$R = A D A^T$$

where D is diagonal and A is orthogonal.

- Eigenvectors of Covariance/Correlation matrix are PC's
Columns of A are PC's
- Diagonal entries of D (=eigenvalues) are variances along PC's (usually sorted in decreasing order)
- R: Function “princomp”

PCA (Version 4): Singular Value Decomposition

- **Singular Value Decomposition:**

Every R can be rewritten as

$$R = U D V^T$$

where D is diagonal and U, V are orthogonal.

- Columns of V are PC's
- Diagonal entries of D are “singular values”; related to standard deviation along PC's (usually sorted in decreasing order)
- UD contains samples measured in PC coordinates
- R : Function “prcomp”

Example: Headsize of sons

Standard deviation in direction of 1.PC,
 $\text{Var} = 12.69^2 = 167.77$

```
> summary(head_pca, loadings = TRUE)
Importance of components:
```

	Comp.1	Comp.2
standard deviation	12.69	5.22
Proportion of Variance	0.86	0.14
Cumulative Proportion	0.86	1.00

Standard deviation in direction of 2.PC,
 $\text{Var} = 5.22^2 = 28.33$
 Total Variance = $167.77 + 28.33 = 196.1$

Loadings:

	Comp.1	Comp.2
head1	0.69	-0.72
head2	0.72	0.69

1.PC contains
 $167.77/196.1 = 0.86$
 of total variance

2.PC contains
 $28.33/196.1 = 0.14$
 of total variance

$$y_1 = 0.69 \cdot x_1 + 0.72 \cdot x_2$$

$$y_2 = -0.72 \cdot x_1 + 0.69 \cdot x_2$$

Computing PC scores

- Subtract mean of all variables
- Output of princomp: \$scores
First column corresponds to coordinate in direction of 1.PC,
Second col. corresponds to coordinate in direction of 2.PC,
etc.
- Manually (e.g. for new observations):
Scalar product of loading of i^{th} PC gives coordinate in
direction of i^{th} PC
- Predict new scores: Use function “predict”
(see ?predict.princomp)
- Example: Headsize of sons

Interpretation of PCs

- Oftentimes hard
- Look at loadings and try to interpret:

Loadings :

	Comp. 1	Comp. 2
head1	0.69	-0.72
head2	0.72	0.69

Difference in head sizes
of both sons

Average head size of both sons

To scale or not to scale...

- R: In princomp, option “cor = TRUE” scales variables
Alternatively: Use correlation matrix instead of covariance matrix
- Use correlation, if different units are compared
- Using covariance will find the variable with largest spread as 1. PC
- Example: Blood Measurement

How many PC's?

- No clear cut rules, only rules of thumb
- **Rule of thumb 1:** Cumulative proportion should be at least 0.8 (i.e. 80% of variance is captured)
- **Rule of thumb 2:** Keep only PC's with above-average variance
(if correlation matrix / scaled data was used, this implies: keep only PC's with eigenvalues at least one)
- **Rule of thumb 3:** Look at scree plot; keep only PC's before the “elbow” (if there is any...)

How many PC's: Blood Example

Rule 1: 5 PC's


Importance of components:

	Comp.1	Comp.2	Comp.3	Comp.4	Comp.5	Comp.6
Standard deviation	1.6710100	1.2375848	1.1177138	0.88227419	0.78839505	0.69917350
Proportion of Variance	0.3490343	0.1914520	0.1561605	0.09730097	0.07769584	0.06110545
Cumulative Proportion	0.3490343	0.5404863	0.6966468	0.79394778	0.87164363	0.93274908

	Comp.7	Comp.8
Standard deviation	0.66002394	0.31996216
Proportion of Variance	0.05445395	0.01279697
Cumulative Proportion	0.98720303	1.00000000

Rule 3: Ellbow after PC 1 (?)

blood_pcaacor



Rule 2: 3 PC's

Mathematical example in detail: Computing eigenvalues and eigenvectors

- See blackboard


Case study: Heptathlon Seoul 1988

Biplot: Show info on samples AND variables

Approximately true:

- Data points: Projection on first two PCs
Distance in Biplot \sim True Distance
- Projection of sample onto arrow gives original (scaled) value of that variable
- Arrowlength: Variance of variabel
- Angle between Arrows: Correlation

Approximation is often crude;
good for quick overview



PCA: Eigendecomposition vs. SVD

- PCA based on Eigendecomposition: princomp
 - + easier to understand mathematical background
 - + more convenient summary method
- PCA based on SVD: prcomp
 - + numerically more stable
 - + still works if more dimensions than samples
- Both methods give same results up to small numerical differences

Concepts to know

- 4 definitions of PCA
- Interpretation: Output of princomp, biplot
- Predict scores for new observations
- How many PC's?
- Scale or not?
- Know advantages of PCA based on SVD

R functions to know

- princomp, biplot
- (prcomp – just know that it exists and that it does the SVD approach)