

# Package ‘Rdta’

January 27, 2024

**Version** 1.0.1

**Date** 2024-1-26

**Title** Data Transforming Augmentation for Linear Mixed Models

**Author** Hyungsuk Tak, Kisung You, Sujit K. Ghosh, and Bingyue Su

**Maintainer** Hyungsuk Tak <hyungsuk.tak@gmail.com>

**Depends** R (>= 2.2.0)

**Imports** MCMCpack(>= 1.4-4), mvtnorm(>= 1.0-11), Rdpack, stats

**Description** We provide a toolbox to fit univariate and multivariate linear mixed models via data transforming augmentation. Users can also fit these models via typical data augmentation for a comparison. It returns either maximum likelihood estimates of unknown model parameters (hyperparameters) via an EM algorithm or posterior samples of those parameters via MCMC. Also see Tak et al. (2019) <[doi:10.1080/10618600.2019.1704295](https://doi.org/10.1080/10618600.2019.1704295)>.

**License** GPL-2

**Encoding** UTF-8

**RdMacros** Rdpack

**RoxygenNote** 6.1.1

**NeedsCompilation** no

**Repository** CRAN

**Date/Publication** 2024-01-27 07:30:10 UTC

## R topics documented:

lmm . . . . .	2
Rdta . . . . .	5
<b>Index</b>	<b>8</b>

---

lmm	<i>Fitting univariate and multivariate linear mixed models via data transforming augmentation</i>
-----	---

---

### Description

The function `lmm` fits univariate and multivariate linear mixed models (also called two-level Gaussian hierarchical models) whose first-level hierarchy is about a distribution of observed data and second-level hierarchy is about a prior distribution of random effects.

### Usage

```
lmm(y, v, x = 1, n.burn, n.sample, tol = 1e-10,
    method = "em", dta = TRUE, print.time = FALSE)
```

### Arguments

<code>y</code>	Response variable. In a univariate case, it is a vector of length $k$ for the observed data. In a multivariate case, it is a ( $k$ by $p$ ) matrix, where $k$ is the number of observations and $p$ denotes the dimensionality.
<code>v</code>	Known measurement error variance. In a univariate case, it is a vector of length $k$ . In a multivariate case, it is a ( $p, p, k$ ) array of known measurement error covariance matrices, i.e., each of the $k$ array components is a ( $p$ by $p$ ) covariance matrix.
<code>x</code>	(Optional) Covariate information. If there is one covariate for each object, e.g., weight, it is a vector of length $k$ for the weight. If there are two covariates for each object, e.g., weight and height, it is a ( $k$ by 2) matrix, where each column contains a covariate variable. Default is no covariate ( $x = 1$ ).
<code>n.burn</code>	Number of warming-up iterations for a Markov chain Monte Carlo method. It must be specified for <code>method = "mcmc"</code>
<code>n.sample</code>	Number of iterations (size of a posterior sample for each parameter) for a Markov chain Monte Carlo method. It must be specified for <code>method = "mcmc"</code>
<code>tol</code>	Tolerance that determines the stopping rule of the EM algorithm. The EM algorithm iterates until the change of log-likelihood function is within the tolerance. Default is $1e-10$ .
<code>method</code>	"em" will return maximum likelihood estimates of the unknown hyper-parameters and "mcmc" returns posterior samples of those parameters.
<code>dta</code>	A logical; Data transforming augmentation is used if <code>dta = TRUE</code> , and typical data augmentation is used if <code>dta = FALSE</code> .
<code>print.time</code>	A logical; TRUE to display two time stamps for initiation and termination, FALSE otherwise.

## Details

For each group  $i$ , let  $y_i$  be an unbiased estimate of random effect  $\theta_i$ , and  $V_i$  be a known measurement error variance. The linear mixed model of interest is specified as follows:

$$[y_i | \theta_i] \sim N(\theta_i, V_i)$$

$$[\theta_i | \mu_{0i}, A] \sim N(\mu_{0i}, A)$$

$$\mu_{0i} = x_i' \beta$$

independently for  $i = 1, \dots, k$ , where  $k$  is the number of groups (units) and dimension of each element is appropriately adjusted in a multivariate case.

The function `lmm` produces maximum likelihood estimates of hyper-parameters,  $A$  and  $\beta$ , their update histories of EM iterations, and the number of EM iterations if method is "em".

For a Bayesian implementation, we put a jointly uniform prior distribution on  $A$  and  $\beta$ , i.e.,

$$f(A, \beta) \propto 1,$$

which is known to have good frequency properties. This joint prior distribution is improper, but their resulting posterior distribution is proper if  $k \geq m + p + 2$ , where  $k$  is the number of groups,  $m$  is the number of regression coefficients, and  $p$  is the dimension of  $y_i$ . We note that an R package `Rgbb` also fits this model in a univariate case ( $p = 1$ ) via ADM (approximation for density maximization). `lmm` produces the posterior samples through a Gibbs sampler if method is "bayes".

## Value

The outcome of `lmm` is composed of:

**A** If method is "mcmc". It contains the posterior sample of  $A$ .

**beta** If method is "mcmc". It contains the posterior sample of  $\beta$ .

**A.mle** If method is "em". It contains the maximum likelihood estimate of  $A$ .

**beta.mle** If method is "em". It contains the maximum likelihood estimate of  $\beta$ .

**A.trace** If method is "em". It contains the update history of  $A$  at each iteration.

**beta.trace** If method is "em". It contains the update history of  $\beta$  at each iteration.

**n.iter** If method is "em". It contains the number of EM iterations.

## Author(s)

Hyungsuk Tak (maintainer), Kisung You, Sujit K. Ghosh, and Bingyue Su

## References

Tak, You, Ghosh, Su, Kelly (2019), "Data Transforming Augmentation for Heteroscedastic Models" [doi:10.1080/10618600.2019.1704295](https://doi.org/10.1080/10618600.2019.1704295)

## Examples

```

#### Univariate linear mixed model

# response variable for 10 objects
y <- c(5.42, -1.91, 2.82, -0.14, -1.83, 3.44, 6.18, -1.20, 2.68, 1.12)
# corresponding measurement error standard deviations
se <- c(1.05, 1.15, 1.22, 1.45, 1.30, 1.29, 1.31, 1.10, 1.23, 1.11)
# one covariate information for 10 objects
x <- c(2, 3, 0, 2, 3, 0, 1, 1, 0, 0)

## Fitting without covariate information
# (DTA) maximum likelihood estimates of A and beta via an EM algorithm
res <- lmm(y = y, v = se^2, method = "em", dta = TRUE)
# (DTA) posterior samples of A and beta via an MCMC method
res <- lmm(y = y, v = se^2, n.burn = 1e1, n.sample = 1e1,
          method = "mcmc", dta = TRUE)
# (DA) maximum likelihood estimates of A and beta via an EM algorithm
res <- lmm(y = y, v = se^2, method = "em", dta = FALSE)
# (DA) posterior samples of A and beta via an MCMC method
res <- lmm(y = y, v = se^2, n.burn = 1e1, n.sample = 1e1,
          method = "mcmc", dta = FALSE)

## Fitting with the covariate information
# (DTA) maximum likelihood estimates of A and beta via an EM algorithm
res <- lmm(y = y, v = se^2, x = x, method = "em", dta = TRUE)
# (DTA) posterior samples of A and beta via an MCMC method
res <- lmm(y = y, v = se^2, x = x, n.burn = 1e1, n.sample = 1e1,
          method = "mcmc", dta = TRUE)
# (DA) maximum likelihood estimates of A and beta via an EM algorithm
res <- lmm(y = y, v = se^2, x = x, method = "em", dta = FALSE)
# (DA) posterior samples of A and beta via an MCMC method
res <- lmm(y = y, v = se^2, x = x, n.burn = 1e1, n.sample = 1e1,
          method = "mcmc", dta = FALSE)

#### Multivariate linear mixed model

# (arbitrary) 10 hospital profiling data (two response variables)
y1 <- c(10.19, 11.53, 16.28, 12.32, 12.84, 11.85, 14.81, 13.24, 14.43, 9.35)
y2 <- c(12.06, 14.97, 11.50, 17.88, 19.21, 14.69, 13.96, 11.07, 12.71, 9.63)
y <- cbind(y1, y2)

# making measurement error covariance matrices for 10 hospitals
n <- c(24, 34, 38, 42, 49, 50, 79, 84, 96, 102) # number of patients
v0 <- matrix(c(186.87, 120.43, 120.43, 250.60), nrow = 2) # common cov matrix
temp <- sapply(1 : length(n), function(j) { v0 / n[j] })
v <- array(temp, dim = c(2, 2, length(n)))

# covariate information (severity measure)
severity <- c(0.45, 0.67, 0.46, 0.56, 0.86, 0.24, 0.34, 0.58, 0.35, 0.17)

## Fitting without covariate information

```

```
# (DTA) maximum likelihood estimates of A and beta via an EM algorithm
res <- lmm(y = y, v = v, method = "em", dta = TRUE)

# (DTA) posterior samples of A and beta via an MCMC method
res <- lmm(y = y, v = v, n.burn = 1e1, n.sample = 1e1,
          method = "mcmc", dta = TRUE)

# (DA) maximum likelihood estimates of A and beta via an EM algorithm
res <- lmm(y = y, v = v, method = "em", dta = FALSE)

# (DA) posterior samples of A and beta via an MCMC method
res <- lmm(y = y, v = v, n.burn = 1e1, n.sample = 1e1,
          method = "mcmc", dta = FALSE)

## Fitting with the covariate information
# (DTA) maximum likelihood estimates of A and beta via an EM algorithm
res <- lmm(y = y, v = v, x = severity, method = "em", dta = TRUE)

# (DTA) posterior samples of A and beta via an MCMC method
res <- lmm(y = y, v = v, x = severity, n.burn = 1e1, n.sample = 1e1,
          method = "mcmc", dta = TRUE)

# (DA) maximum likelihood estimates of A and beta via an EM algorithm
res <- lmm(y = y, v = v, x = severity, method = "em", dta = FALSE)

# (DA) posterior samples of A and beta via an MCMC method
res <- lmm(y = y, v = v, x = severity, n.burn = 1e1, n.sample = 1e1,
          method = "mcmc", dta = FALSE)
```

## Description

The R package **Rdta** provides a toolbox to fit univariate and multivariate linear mixed models via data transforming augmentation. Users can also fit these models via typical data augmentation for a comparison. It returns either maximum likelihood estimates of unknown model parameters (hyper-parameters) via an EM algorithm or posterior samples of those parameters via a Markov chain Monte Carlo method.

**Details**

Package: Rdta  
Type: Package  
Version: 1.0.1  
Date: 2024-1-26  
License: GPL-2  
Main functions: [lmm](#)

**Author(s)**

Hyungsuk Tak (maintainer), Kisung You, Sujit K. Ghosh, and Bingyue Su

**References**

Tak, You, Ghosh, Su, Kelly (2019), "Data Transforming Augmentation for Heteroscedastic Models"  
<[doi:10.1080/10618600.2019.1704295](https://doi.org/10.1080/10618600.2019.1704295)>

# Index

lmm, [2](#), [7](#)

Rdta, [5](#)

Rdta-package (Rdta), [5](#)