

# Dealing with missing values – part 2

Applied Multivariate Statistics – Spring 2013




# Overview

- More on Single Imputation: Shortcomings
- Multiple Imputation: Accounting for uncertainty

# Single Imputation

- Unconditional Mean
- Unconditional Distribution
- Conditional Mean
- Conditional Distribution

Easy / Inaccurate



Hard / Accurate

## Example: Blood Pressure - Revisited


- 30 participants in January (X) and February (Y)
- MCAR: Delete 23 Y values randomly
- MAR: Keep Y only where  $X > 140$  (follow-up)
- MNAR: Record Y only where  $Y > 140$  (test everybody again but only keep values of critical participants)

X	Y			
	Complete	MCAR	MAR	MNAR
Data for individual participants				
169	148	148	148	148
126	123	—	—	—
132	149	—	—	149
160	169	—	169	169
105	138	—	—	—
116	102	—	—	—
125	88	—	—	—
112	100	—	—	—
133	150	—	—	150
94	113	—	—	—
109	96	—	—	—
109	78	—	—	—
106	148	—	—	148
176	137	—	137	—
128	155	—	—	155
131	131	—	—	—
130	101	101	—	—
145	155	—	155	155
136	140	—	—	—
146	134	—	134	—
111	129	—	—	—
97	85	85	—	—
134	124	124	—	—
153	112	—	112	—
118	118	—	—	—
137	122	122	—	—
101	119	—	—	—
103	106	106	—	—
78	74	74	—	—
151	113	—	113	—


# Example: Blood Pressure

Black points are missing (MAR)

True values



MAR




# Unconditional Mean


+ Mean of Y ok

- Variance of Y wrong

True values



Unconditional Mean




# Unconditional Distribution


+ Mean of Y ok, Variance better

- Correlation btw X and Y wrong

True values




Unconditional Distribution




# Conditional Mean

- + Conditional Mean of Y ok
- + Correlation ok
- (Conditional) Variance wrong

True values



Conditional Mean







# Conditional Distribution

- + Conditional Mean of Y ok
- + Correlation ok
- + Conditional Variance of Y ok

**True values**





**Conditional Distribution**



# Problem: We ignore uncertainty

True values


Conditional Distribution



# Problem of Single Imputation

- Too optimistic: Imputation model (e.g. in  $Y = a + bX$ ) is **just estimated**, but not the true model
- Thus, imputed values have some uncertainty
- Single Imputation ignores this uncertainty
- Coverage probability of confidence intervals is wrong
  
- Solution: Multiple Imputation
  - Incorporates both
    - residual error
    - model uncertainty (excluding model mis-specification)


# Multiple Imputation: Idea



# Multiple Imputation: Idea


- Need special imputation schemes that include both
  - uncertainty of residuals
  - uncertainty of model  
(e.g. values of intercept  $a$  and slope  $b$ )
- Rough idea:
  - Fill in random values
  - Iteratively predict values for each variable until some convergence is reached (as in missForest)
  - Sample values for residuals **AND for  $(a,b)$**
- Gibbs sampler is used
- Excellent for intuition (by one of the big guys in the field):  
<http://sites.stat.psu.edu/~jls/mifaq.html>

# Multiple Imputation: Intuition




- Predict missing values accounting for
- Uncertainty of residuals
  - Uncertainty of parameter estimates

# Multiple Imputation: Intuition



- Predict missing values accounting for
- Uncertainty of residuals
  - Uncertainty of parameter estimates


# Multiple Imputation: Intuition



- Predict missing values accounting for
- Uncertainty of residuals
  - Uncertainty of parameter estimates




# Multiple Imputation: Intuition




- Predict missing values accounting for
- Uncertainty of residuals
  - Uncertainty of parameter estimates

# Multiple Imputation: Intuition



- Predict missing values accounting for
- Uncertainty of residuals
  - Uncertainty of parameter estimates

# Multiple Imputation: Intuition




Predict missing values accounting for

- Uncertainty of residuals
- Uncertainty of parameter estimates

## Multiple Imputation: Gibbs sampler (Not for exam)

- Iteration  $t$ ; repeat until convergence:  
For each variable  $i$ :



## R package: MICE

### Multiple Imputation with Chained Equations

- MICE has good default settings; don't worry about the data type
- Defaults for data types of columns:
  - **numeric**: Predictive Mean Matching (pmm)  
(like fancy linear regression; faster alternative: linear regression)
  - **factor, 2 lev**: Logistic Regression (logreg)
  - **factor, >2 lev**: Multinomial logit model (polyreg)
  - **ordered, >2 lev**: Ordered logit model (polr)


# Aggregation of estimates

- $\hat{Q}_i$  : Estimate of imputation  $i$   
 $U_i$  : Variance of estimate (= square of std. error)
- Assume:  $\frac{\hat{Q} - Q}{\sqrt{\bar{U}}} \approx N(0, 1)$
- Average estimate:  $\bar{Q} = \frac{1}{m} \sum_{j=1}^m \hat{Q}_j$ 

Here was a typo
- Within-imputation variance:  $\bar{U} = \frac{1}{m} \sum_{j=1}^m \hat{U}_j$
- Between-imputation variance:  $B = \frac{1}{m-1} \sum_{j=1}^m (\hat{Q}_j - \bar{Q})^2$
- Total variance:  $T = \bar{U} + (1 + \frac{1}{m})B$
- Approximately:  $\frac{\bar{Q} - Q}{\sqrt{T}} \sim t_\nu$  with  $\nu = (m - 1) \left(1 + \frac{m\bar{U}}{(1+m)B}\right)^2$
- 95%-CI:  $\bar{Q} \pm t_{\nu; 0.975} \sqrt{T}$

# Multiple Imputation with MICE

Do manually, if you have non standard analysis



# How much uncertainty due to missings?

- Relative increase in variance due to nonresponse:

$$r = \frac{(1 + \frac{1}{m})B}{U}$$

- Fraction (or rate) of missing information fmi:  
(!! Not the same as fraction of missing OBSERVATIONS)

$$fmi = \frac{r + \frac{2}{\nu+3}}{r+1}$$

- Proportion of the total variance that is attributed to the missing data:

$$\lambda = \frac{B(1 + \frac{1}{m})}{T}$$

Returned by mice





Rule of thumb:

- Preliminary analysis:  $m = 5$
- Paper:  $m = 20$  or even  $m = 50$

## How many imputations?

- Surprisingly few!
- Efficiency compared to  $m = \infty$  depends on fmi:

$$eff = \left(1 + \frac{fmi}{m}\right)^{-1}$$

- Examples (eff in %):

Oftentimes OK

Perfect !

M	fmi=0.1	fmi=0.3	fmi=0.5	fmi=0.7	fmi=0.9
3	97	91	86	81	77
5	98	94	91	88	85
10	99	97	95	93	92
20	100	99	98	97	96

# Concepts to know

- Idea of mice
- How to aggregate results from imputed data sets?
- How many imputations?

# R functions to know

- mice, with, pool

# Next time

- Multidimensional Scaling
- Distance metrics